



[Web](#) [Images](#) [Video](#) [News](#) [Maps](#) [more »](#)

man glcnac antibody thiol OR Cysteine iodoacetamide

Search

[Advanced Scholar Search](#)

[Scholar Preferences](#)

[Scholar Help](#)

Scholar [All articles](#) - [Recent articles](#) Results **1 - 10** of about **227** for **man glcnac antibody thiol OR Cysteine iodoacetamide** with **Safesearch on**. (0.29 seconds)

[Mass Spectrometry of the Humanized Monoclonal **Antibody** Campath 1H - all 3 versions »](#)

DS Ashton, OR Beddell, DJ Cooper, SJ Craig, AC ... - Analytical Chemistry, 1995 - pubs.acs.org

... DL-**Cysteine**, dithiothreitol ... variant of bacterial subtilisin and a bacterial **thiol**

oxidoreductase were ... Chester, UK A solution of CAMPATH 1H **antibody** in phosphate ...

[Cited by 18](#) - [Related Articles](#) - [Web Search](#)

[\[CITATION\] Plant-derived mouse IgG monoclonal **antibody** fused to KDEL endoplasmic reticulum-retention signal is ... - all 2 versions »](#)

A Triguero, G Cabrera, JA Cremata, CT Yuen, J ... - Plant Biotechnology Journal, 2005 - Blackwell Synergy

... C with anti- β 1,2-Xyl and anti- α 1,3-Fuc **antibodies** (diluted 1 ... Relative retention times

were calculated, referred to that of the standard **Man 3 GlcNAc 2-2AB** ...

[Cited by 18](#) - [Related Articles](#) - [Web Search](#)

[Impact of variable domain glycosylation on **antibody** clearance: An LC/MS characterization - all 3 versions »](#)

L Huang, S Bioisi, KR Bales, U Kuchibhotla - Analytical Biochemistry, 2006 - Elsevier

... Another **GlcNAc** must be linked to only available HexNAc ... its linkage to GlcNAc β (1 →

2)**Man** could not ... captured using immobilized goat anti-human Fc γ **antibody**. ...

[Cited by 11](#) - [Related Articles](#) - [Web Search](#)

[\[CITATION\] Structural Characterization of a Recombinant **Antibody** Produced in Yeast](#)

PC Gough, IA Kadura, GM Kelly, A Kreuzman, J Larson ...

[Related Articles](#) - [Web Search](#)

[Biochemical and functional characterization of the Tn-specific lectin from Salvia sclarea seeds - all 4 versions »](#)

A Medeiros, S Bianchi, JJ Calvete, H Balter, S Bay ... - FEBS Journal, 2000 - FEBS

... Anti-Tn monoclonal **antibody** 83D4 (IgM,) [22] was ... molar excess of **iodoacetamide** over

cysteine residues; and ... chains of SSL are composed of **GlcNAc**, **Man**, Xyl, and ...

[Cited by 17](#) - [Related Articles](#) - [Web Search](#) - [BL Direct](#)

[N-Glycosylation of a mouse IgG expressed in transgenic tobacco plants - all 3 versions »](#)

M Cabanes-Macheteau - Glycobiology, 1999 - Soc Glycobiology

... **GlcNAc 2** core substituted by terminal **GlcNAc** residues ... **Man-8**), and soybean agglutinin

(**Man-9**) as ... to pulsed amperometric detection; MAb, monoclonal **antibody**; 1 H ...

[Cited by 118](#) - [Related Articles](#) - [Web Search](#) - [BL Direct](#)

[... Specific Glycosylation of an Aglycosylated Human IgG1-Fc **Antibody** Protein Generates Neoglycoproteins ... - all 11 versions »](#)

GM Watt, J Lund, M Levens, VSK Kolli, R Jefferis, ... - Chemistry & Biology, 2003 - Elsevier

... 1). Epitope mapping using an extensive panel of monoclonal **antibodies** directed

against the human IgG1 **antibody** revealed that ... [-S(**GlcNAc**) 2 **Man**] 15 25,030 ...

[Cited by 11](#) - [Related Articles](#) - [Web Search](#)

[Glycosylation of the variable region of immunoglobulin G—site specific maturation of the sugar ...](#) - [all 4 versions »](#)

T Endo, A Wright, SL Morrison, A Kobata - Molecular Immunology, 1995 - Elsevier

... Purified secreted **antibody** from each transfectoma ... M, EDTA to 2 mM and **cysteine** to 10 ... **Man**?1 + 4GlcNAcBI+ 4GlcNAcOT (**Man** - **GlcNAc** - **GlcNAc**,,) was prepared from ...

[Cited by 15](#) - [Related Articles](#) - [Web Search](#) - [BL Direct](#)

[The cattle tick antigen, Bm95, expressed in Pichia pastoris contains short chains of N-and O-glycans](#) - [all 4 versions »](#)

LJ González, JA Cremata, Y Guanche, Y Ramos, A ... - Archives of Biochemistry and Biophysics, 2004 - Elsevier

... 7]. However, B. microplus isolations show low susceptibility to **antibodies** developed

in ... a database of the retention times relative to **Man** 7 **GlcNAc** 2 -4ABA ...

[Cited by 3](#) - [Related Articles](#) - [Web Search](#)

[Mass spectrometric analysis reveals a **cysteine** bridge between residues 2 and 61 of the auxin-binding ...](#) - [all 3 versions »](#)

C Feckler, G Muster, W Feser, A Römer, K Palme - FEBS Letters, 2001 - Elsevier

... were based on epitope mapping using monoclonal **antibodies** against peptides ... type sugar

predicted from earlier results was considered (**GlcNAc** 2 **Man** 9 , 1865.6 ...

[Cited by 4](#) - [Related Articles](#) - [Web Search](#)

Key authors: [D Williams](#) - [A Helenius](#) - [A Parodi](#) - [E Trombetta](#) - [B Davis](#)

Google ►

Result Page: [1](#) [2](#) [3](#) [4](#) [5](#) [6](#) [7](#) [8](#) [9](#) [10](#) [Next](#)

man glcnac antibody thiol OR Cyste

Search

[Google Home](#) - [About Google](#) - [About Google Scholar](#)

©2008 Google